

On the complexity of finding a sun in a graph

Chính T. Hoàng*

Abstract

The sun is the graph obtained from a cycle of length even and at least six by adding edges to make the even-indexed vertices pairwise adjacent. Suns play an important role in the study of strongly chordal graphs. A graph is chordal if it does not contain an induced cycle of length at least four. A graph is strongly chordal if it is chordal and every even cycle has a chord joining vertices whose distance on the cycle is odd. Farber proved that a graph is strongly chordal if and only if it is chordal and contains no induced suns. There are well known polynomial-time algorithms for recognizing a sun in a chordal graph. Recently, polynomial-time algorithms for finding a sun for a larger class of graphs, the so-called HHD-free graphs, have been discovered. In this paper, we prove the problem of deciding whether an arbitrary graph contains a sun in NP-complete.

Keywords: chordal graph, strongly chordal graph, sun

1 Introduction

A *hole* is an induced cycle with at least four vertices. A graph is *chordal* if it does not contain a hole as an induced subgraph. Farber [6] defined a graph to be *strongly chordal* if it is chordal and every cycle in the graph on $2k$ vertices, $k \geq 3$, has a chord uv such that each segment of the cycle from u to v has an odd number of edges. We denote by k -*sun* the graph obtained from a cycle of length $2k$ ($k \geq 3$) by adding edges to make the even-indexed vertices pairwise adjacent. Figure 1 shows a 5-sun. A *sun* is simply a k -sun for some $k \geq 3$. Farber showed [6] that a graph is strongly chordal if and only if it is chordal and does not contain a sun as induced subgraph. Farber's motivation was a polynomial-time algorithm for the minimum weighted dominating set problem for strongly chordal graphs. The problem is NP-hard for chordal graphs [1]. In this paper, we prove that it is NP-hard to find a sun in an arbitrary graph. This result is motivated by the following discussion on chordal and strongly chordal graphs. For more information on this topics, see [3, 7].

We use $N(x)$ to denote the set of vertices adjacent to vertex x in a graph G . Define $N[x] = N(x) \cup \{x\}$. A vertex x in a graph is *simplicial* if $N(x)$ induces a complete graph. It is well known [4] that graph G is chordal if and only if every induced subgraph H of G contains a simplicial vertex of H . Farber proved [6] an analogous characterization for strongly chordal graphs. A vertex x in a

*Department of Physics and Computer Science, Wilfrid Laurier University, Canada. Research supported by NSERC.
email: choang@wlu.ca




Figure 1: The 5-sun




Figure 2: The house, the hole and the domino

graph is *simple* if the vertices in $N(x)$ can be ordered as $x_1, x_2, \dots, x_k$ such that $N[x_1] \subseteq N[x_2] \subseteq \dots \subseteq N[x_k]$. Thus, every simple vertex is simplicial. For a graph G , let $\mathcal{R} = v_1, v_2, \dots, v_n$ be an ordering of vertices of G . Let $G(i) = G[\{v_i, v_{i+1}, \dots, v_n\}]$, i.e., the subgraph induced in G by the set v_i through v_n of vertices. $\mathcal{R}$ is a *simple elimination ordering* for G if v_i is simple in $G(i)$, $1 \leq i \leq n$. The following is due to Farber [6]:

Theorem 1 ([6]) *The following are equivalent for any graph G :*

- G is strongly chordal.
- G is chordal and does not contain a sun.
- Vertices of G admit a simple elimination ordering.

Thus, suns play an important role in the studies of chordal and strongly chordal graphs. There are well known algorithms [17, 12] to test whether a chordal graph is strongly chordal and thus whether it contains a sun. It is natural to investigate the problem for larger classes of graphs. A graph is HHD-free if it does not contain a house, a hole, or a domino (see figure 2). Every chordal graph is a HHD-free graph. HHD-free graphs [10] have several properties analogous to those of chordal graphs. Brandstädt [2] proposed the problem of finding a sun in a HHD-free graph. This problem was proved to be polynomial-time solvable in [13] and [5]. In this paper, we will prove the following

Theorem 2 *It is NP-complete to decide whether a graph contains a sun.*

Denote by k -hole the hole on k vertices. A k -antihole is the complement of a k -hole. A graph is *weakly chordal* [8] if it does not contain a k -hole or k -antihole with $k \geq 5$. Weakly chordal graphs generalize chordal graphs in a natural way, and they are known to be perfect and have many interesting algorithmic properties (see [9]). In spite of Theorem 2, it is conceivable there are polynomial-time algorithms to solve the sun recognition problem for weakly chordal graphs or even perfect graphs [15]. In this spirit, we will refine Theorem 2 to obtain a stronger result.

Theorem 3 *It is NP-complete to decide whether a graph G contains a sun, even when G does not contain a k -antihole with $k \geq 7$.*

Let k -CLIQUE (respectively, k -SUN) be the problem whose instance is a graph G and an integer k , for which the question to be answered is whether G contains a clique on k vertices (respectively, k -SUN). It is well known [11] that k -CLIQUE is NP-complete. It is not difficult to prove, but perhaps interesting to note that k -SUN is also NP-complete. Observe that if k is a constant (not part of the input), then the two problems can obviously be solved in polynomial time.

Theorem 4 *k -SUN is NP-complete.*

Note that Theorem 2 implies Theorem 4: To decide whether a graph contains a sun, we only need solve $O(n)$ instances of k -SUN with k running from 3 to $n/2$, where n is the number of vertices of the graph. However, we have a short and direct proof of Theorem 4. We will give the proofs of Theorems 2, 3 and 4 in the remainder of the paper.

2 The proofs

First, we need introduce some definitions. For simplicity, we will say a vertex x *sees* a vertex y if x is adjacent to y ; otherwise, we will say x *misses* y . Let G, F be two vertex-disjoint graphs and let x be a vertex of G . We say that a graph H is obtained from G by *substituting* F for x if H is obtained by replacing x by F in G and adding the edge ab for any $a \in V(G) - \{x\}$, and any $b \in F$ whenever ax is an edge of G . In the proofs, we will often use the observation that every vertex in $H - F$ either sees all, or misses all, vertices of F .

By $(c_1, c_2, \dots, c_k, e_1, e_2, \dots, e_k)$ we denote the k -sun with vertices $c_1, c_2, \dots, c_k, e_1, e_2, \dots, e_k$ such that $c_1, c_2, \dots, c_k$ induce a clique, $e_1, e_2, \dots, e_k$ induce a stable set, each e_i has degree two and sees c_i, c_{i+1} with the subscripts taken modulo k . The vertices e_i will be called the *ears* of the k -sun. A *triangle* is a clique on three vertices.

We will rely on the following NP-complete problem due to Poljak [14].

STABLE SET IN TRIANGLE-FREE GRAPHS

Instance: A triangle-free graph G , an integer k .

Question: Does G contain a stable set with k vertices?

Proof of Theorem 2. We will reduce STABLE SET IN TRIANGLE-FREE GRAPHS to the problem of finding a sun in a graph.




Figure 3: The graph G




Figure 4: The graph $f(G,4)$

Let $G = (V, E)$ be a triangle-free graph with $V = \{v_1, v_2, \dots, v_n\}$, and without loss of generality assume $k \geq 4$. Define a graph $f(G, k)$ from G as follows. Substitute for each vertex v_i a clique $V_i = \{v_i^1, v_i^2, \dots, v_i^k\}$; add a clique W with vertices $u_1, w_1, \dots, u_k, w_k$; add a stable set X with vertices $x_1, \dots, x_k$; for $i = 1, 2, \dots, k$, add edges $x_i w_i$ and $x_i u_{i+1}$ (the subscripts are taken module k); for $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, k$, add edges $v_i^j u_j, v_i^j w_j$. Figure 3 shows a graph G whose graph $f(G, 4)$ is shown in Figure 4 (for clarity, we do not show all edges of $f(G, 4)$; all adjacency between V_1 and W , and between V_2 and W are shown, adjacency between V_3 and W are not shown; the thick line between V_1 and V_2 (and between V_2 and V_3) represents all possible edges between the two sets; there are no edges between V_1 and V_3 ; each of the sets V_i , W induces a clique; the set X induces a stable set.) We will often rely on the following observations.

Observation 1 Suppose G is triangle-free. Then $f(G, k)$ does not contain a triangle each of whose vertices belongs to a distinct V_i . $\square$

Observation 2 Let x be a vertex in V_i , y be a vertex in V_j with $i \neq j$. If x and y have a common neighbor z in W , then $N(x) \cap W = N(y) \cap W$. $\square$

The theorem follows from the following claim.

Claim 1 G has a stable set with k vertices if and only if $f(G, k)$ contains a sun.

Proof of Claim 1. Suppose G has a stable set with vertices $v_1, v_2, \dots, v_k$. Then $f(G, k)$ has a $2k$ -sun $(c_1, c_2, \dots, c_{2k}, e_1, e_2, \dots, e_{2k})$ with $e_{2i-1} = v_i^i$, $e_{2i} = x_i$, $c_{2i-1} = u_i$, $c_{2i} = w_i$, for $i = 1, 2, \dots, k$.

Now, suppose $f(G, k)$ contains a sun. Write $U = V_1 \cup V_2 \cup \dots \cup V_n$. We will establish that

Any sun S of $f(G, k)$ is a $2k$ -sun with k ears in $V_1 \cup \dots \cup V_n$. (1)

Consider a sun $S = (c_1, c_2, \dots, c_t, e_1, e_2, \dots, e_t)$ of $f(G, k)$. First, we claim that (with the subscript taken modulo k)

If an ear e_j lies in X , then e_{j-1}, e_j lie in U . (2)

Let x_i be a vertex in X that is an ear e_j of S . We may assume that $c_j = w_i$ and $c_{j+1} = u_{i+1}$. Since e_{j-1} sees w_i and misses u_{i+1} , we have $e_{j-1} \in V_s$ for some s . Similarly, we have $e_{j+1} \in V_r$ for some r . Note that $r \neq s$. So, (2) holds.

Since W is a clique, S must have an ear in $U \cup X$. (2) implies that

U contains an ear of S . (3)

Next, we will prove

If $e_i \in V_j$ then $c_i, c_{i+1} \in W$. (4)

Suppose (4) is false. For simplicity, we may assume $i = 1$ and $j = 1$ (we can always rename the vertices of $f(G, k)$ and S so that this is the case). We will often implicitly use the fact that a vertex in V_a either sees all, or misses all, vertices of V_b whenever $a \neq b$. We will distinguish among several cases.

Case 1: $c_1, c_2 \in V_1$. Since c_3 sees c_1, c_2 and misses e_1 , c_3 cannot be in U . Thus, c_3 is in W . But no vertex in W can see two vertices in V_1 , a contradiction.

Case 2: $c_1 \in V_1, c_2 \in V_j$ for some $j \neq 1$. We may write $j = 2$. Since c_3 (respectively, e_t) sees c_1 and misses e_1 , c_3 (respectively, e_t) cannot be in U . Thus, c_3 and e_t are in W . Observation 2, with $z = c_3, x = c_1, y = c_2$, implies e_t sees c_2 , a contradiction to the definition of S .

Case 3: $c_1 \in V_1, c_2 \in W$. This case is not possible since a vertex in W can have at most one neighbor in any V_j .

Case 4: $c_1, c_2 \in V_j$ for some $j \neq 1$. We may write $j = 2$. Since e_2 sees c_2 and misses c_1 , e_2 is in W . Since e_t sees c_1 and misses c_2 , e_t is in W . But then e_2 sees e_t , a contradiction.

Case 5: $c_1 \in V_j, c_2 \in V_r$ with $j \neq r, j \neq 1, r \neq 1$. In this case, e_1, c_1, c_2 contradict Observation 1.

Case 6: $c_1 \in V_j$ for some $j \neq 1$ and $c_2 \in W$. We may let $j = 2$. If $c_3 \in W$, then Observation 2, with $z = c_2, x = e_1, y = c_1$ implies c_3 sees e_1 , a contradiction to the definition of S . So, we have $c_3 \in U$. Since c_3 misses e_1 , we have $c_3 \notin V_1 \cup V_2$. So, we may assume $c_3 \in V_3$. We have $e_2 \notin W$; for otherwise Observation 2, with $z = c_2, x = c_1, y = c_3$, implies e_2 sees c_1 , a contradiction to the definition of S . We have $e_2 \notin V_1 \cup V_2 \cup V_3$ since e_2 misses e_1 and c_1 . So, we may assume $e_2 \in V_4$. Since e_3 (respectively, c_4 , if it exists) sees c_3 and misses e_1 , Observation 2, with $z = c_2, x = e_1, y = c_3$, implies $e_3 \notin W$ (respectively, $c_4 \notin W$). Since e_3 (respectively, c_4 , if it exists) misses e_1 and e_2 , we have $e_3 \notin V_1 \cup V_2 \cup V_3 \cup V_4$ (respectively, $c_4 \notin V_1 \cup V_2 \cup V_3 \cup V_4$). Now, if $t = 3$, then the three vertices e_3, c_1, c_3 contradict Observation 1. But if $t > 3$, then the three vertices c_4, c_1, c_3 contradict Observation 1.

So (4) holds. Next, we will establish two more assertions (where the subscripts are taken modulo k) below.

$$\text{If an ear } e_j \text{ lies in } U \text{ then } e_{j-1}, e_{j+1} \text{ lie in } X. \quad (5)$$

By (4) and the definition of $f(G, k)$, we may assume $c_i = u_i, c_{i+1} = w_i$. Since x_i is the only vertex of $f(G, k)$ that sees w_i and misses u_i , we have $x_i = e_{j+1}$. Similarly, we have $x_{i-1} = e_{j-1}$. So, (5) holds.

$$\text{If some vertex } x_i \in X \text{ is an ear of } S, \text{ then } x_{i+1} \text{ is also an ear of } S. \quad (6)$$

Let x_i be a vertex in X that is an ear e_j of S . We may assume that $c_j = w_i$ and $c_{j+1} = u_{i+1}$. By (2), we have $e_{j+1} \in V_a$ for some a . By (4), we have $c_{j+2} = w_{i+1}$. By (5), e_{j+2} lies in X , and so we have $e_{j+2} = x_{i+1}$. Thus, (6) holds.

We are now in position to prove (1). From (3), we may assume e_1 lies in U . By (5), we have $e_2 \in X$. By (6), all x_j are ears of S for $j = 1, 2, \dots, k$. It follows from (2) that S has exactly k ears in U . Therefore, S is a $2k$ -sun. We have proved (1).

We continue with the proof of the Claim (and the Theorem). Consider the k ears of S that belong to U . Since each V_i is a clique, it contains at most one ear. So, there are k sets V_i containing an ear of S . Let these sets be $V_1, V_2, \dots, V_k$. Clearly, in G , the vertices $v_1, v_2, \dots, v_k$ form a stable set. $\square$

Proof of Theorem 3. We will use the notation defined in the proof of Theorem 2 with G being a triangle-free graph. We only need prove the graph $f(G, k)$ does not contain a t -antihole with $t \geq 7$. We will prove by contradiction. Suppose $f(G, k)$ contains a t -antihole A with vertices $a_1, a_2, \dots, a_t$ with $t \geq 7$ such that a_i misses a_{i+1} with the subscripts taken modulo k . Since the vertices in X have degree two, none of them can belong to A . Since each V_i is a clique,

$$\text{no two consecutive vertices of } A \text{ can belong to the same } V_i. \quad (7)$$

Similarly,

$$\text{no two consecutive vertices of } A \text{ can belong to } W. \quad (8)$$

Now, we claim that

$$\text{one of } a_i, a_{i+1} \text{ must lie in } W \text{ for all } i. \quad (9)$$

Suppose (9) is false for a_i . For simplicity, we may assume $i = 1$, and so we have $a_1, a_2 \in U$. By (7), we may assume $a_1 \in V_1, a_2 \in V_2$. Clearly, we have $a_t \notin V_1$.

Suppose $a_t \in V_2$. Then a_3 has to be in W , for otherwise a_3 lies in some V_j and so it misses a_t (since it misses a_2) implying $t = 4$, a contradiction. By symmetry, we have $a_{t-1} \in W$. Since a_1 sees a_3 , and a_{t-1} is a common neighbour of a_1 and a_2 , Observation 2 implies that a_2 sees a_3 , a contradiction to the definition of A . So, we have $a_t \notin V_2$.

Suppose $a_t \in W$. By (8), we have $a_{t-1} \in V_j$. If $j = 2$ then a_1 misses a_{t-1} , a contradiction to the definition of A . If $j = 1$ then a_2 misses a_{t-1} implying $t = 4$, a contradiction. So, we may assume $a_{t-1} \in V_3$. Let $j \in \{t-2, t-3\}$. If $a_j \in W$ then since a_2 sees a_t , Observation 2 with $z = a_j, x = a_2, y = a_1$ implies a_1 sees a_t , a contradiction to the definition of A . So, we have $a_{t-2} \in V_m$ for some m , and $a_{t-3} \in V_p$ for some p . Since a_{t-2} misses a_{t-1} , we have $a_{t-2} \notin V_1 \cup V_2 \cup V_3$. So, we may assume $m = 4$. We have $a_{t-3} \in V_2 \cup V_3$, for otherwise the three vertices a_{t-3}, a_{t-1}, a_2 contradict Observation 1. Since a_{t-2} sees a_2 , a_{t-2} sees all of V_2 . Thus, we have $a_{t-3} \notin V_2$, and so $a_{t-3} \in V_3$. Since $t \geq 7$, the vertex a_{t-4} exists. Since a_{t-4} misses a_{t-3} but sees a_{t-1} , a_{t-4} is not in U ; so we have $a_{t-4} \in W$. Observation 2 with $z = a_{t-4}, x = a_{t-1}, y = a_{t-2}$ implies a_{t-1} sees a_t , a contradiction to the definition of A .

Thus, a_t belongs to some V_j which is distinct from V_1, V_2 . It follows from symmetry and the definition of $f(G, k)$ that a_3, a_{t-1} also belong to distinct V_i . Now, the three vertices a_{t-1}, a_1, a_3 contradict Observation 1. So, (9) holds.

From (8) and (9), we may assume without loss of generality that $a_i \in U$ whenever i is odd, and $a_i \in W$ whenever i is even. In particular, t is even and at least eight. The definition of A implies that a_1 sees a_4, a_6 . Thus, we have $\{a_4, a_6\} = \{u_i, w_i\}$ for some i . The definition of $f(G, k)$ means that every vertex of U either sees both a_4, a_6 or misses both of them. But a_3 misses a_4 and sees a_6 , a contradiction. $\square$

Proof of Theorem 4. We will reduce k -CLIQUE to k -SUN. Let G, k be an instance of k -CLIQUE. We may assume $k \geq 4$. Construct a graph $h(G)$ from G by adding a vertex $v(a, b)$ for each edge ab of G , and joining $v(a, b)$ to a and b by an edge of $h(G)$. Let Y be the set of vertices $v(a, b)$. It is easy to see that if G has a clique K on k vertices then $h(G)$ has a k -sun induced by K and some k vertices in Y . If $h(G)$ has a k -sun $(c_1, \dots, c_k, e_1, \dots, e_k)$, then since the vertices in Y have degree two, none of them can be a vertex c_i ; thus, the vertices $c_1, \dots, c_k$ induce a clique on k vertices in G . $\square$

References

- [1] K. S. Booth, and J. H. Johnson, Dominating set in chordal graphs. *SIAM J. Comput.* **11**(1), (1980) 191-199.
- [2] A. Brandstädt, in *Problem Session*, Dagstuhl seminar on robust and approximative algorithms for particular graph classes, Seminar No. 04221, Wadern, Germany, May 2004.
- [3] A. Brandstädt, V.B. Le, and J.P. Spinrad, *Graph Classes: A survey*, SIAM Monographs on Discrete Mathematics and Applications, 1999.

- [4] G.A. Dirac, On rigid circuit graphs, *Abh. Math. Sem. Univ. Hamburg* **25**, (1961) 71–76.
- [5] E. M. Eschen, C. T. Hoàng, and R. Sritharan, An $O(n^3)$ recognition algorithm for hhds-free graphs, *Graphs and Combinatorics* **23**, (2007) Supplement 1: 209–231.
- [6] M. Farber, Characterizations of strongly chordal graphs, *Discrete Mathematics* **43**, (1983) 173–189.
- [7] M. C. Golumbic, *Algorithmic graph theory and perfect graphs*, Academic Press, New York, 1980.
- [8] R. B. Hayward, Weakly Triangulated Graphs, *J. Combin. Theory Ser. B*, **39**, (1985) 200–209.
- [9] R. B. Hayward, J. Spinrad and R. Sritharan, Improved algorithms for weakly chordal graphs. *ACM Transactions on Algorithms* **3**(2), (2007), article 14.
- [10] C. T. Hoàng and N. Khouzam. On brittle graphs. *J. Graph Theory* **12** (1988) 391–404.
- [11] R. Karp. Reducibility Among Combinatorial Problems, in: *Complexity of Computer Computations*, R. E. Miller and J. W. Thatcher (eds), Plenum Press, (1972) 85–103.
- [12] A. Lubiw, Doubly lexical ordering of matrices, *SIAM Journal on Computing* **16**, (1987) 854–879.
- [13] S.D. Nikolopoulos and L. Palios, Recognizing hhds-free graphs, in *Proceedings of the 31st International Workshop on Graph Theoretic Concepts in Computer Science (WG 2005)*, Metz, France, 2005.
- [14] S. Poljak. A note on stable sets and coloring of graphs. *Comment. Math. Univ. Carolinae* **15** (1974), pp. 307–309.
- [15] J. L. Ramrez-Alfonso and B. A. Reed (eds), *Perfect Graphs*, Wiley, 2001.
- [16] D.J. Rose, R.E. Tarjan, and G.S. Leuker, Algorithmic aspects of vertex elimination on graphs, *SIAM Journal on Computing* **5**, (1976) 266–283.
- [17] R. Paige and R. E. Tarjan, Three Partition Refinement Algorithms, *SIAM J. Comput.* **16** (1987) 973–989.